

Please read the instructions carefully before using.

# イブクイック®頭痛薬

頭痛・熱に 第②類医薬品

※ この医薬品は「使用上の注意」をよく読んでお使いください。

アレルギー体質の方は、必ず薬剤師、登録販売者にご相談ください。

Pain Reliever/Fever Reducer  
Ibuprofen combination product



EVE QUICK FOR HEADACHE

Designated Class 2 OTC drug

- EVE QUICK FOR HEADACHE is a formulation designed to work well on headache. It combines ibuprofen, plus allylisopropylacetylurea and anhydrous caffeine which increase its analgesic effects. In addition, it has a blend of magnesium oxide which protects the stomach lining and promotes the absorption of ibuprofen.

<Features>

- It exerts quick and superior effects on the source of headache because its own formulation promotes the absorption of analgesic ingredients.
- It contains ingredients that are gentle on the stomach and protect the stomach lining.

## CAUTIONS FOR USE

### CONTRAINDICATIONS

(Failure to observe the following may cause the present symptoms to worsen or produce adverse drug reactions and/or accidents.)

#### 1. The following people must not take this drug.

- (1) Those with a previous allergic reaction to this drug or ingredients of the drug
- (2) Those who have ever suffered from asthma after taking this drug or other drugs for the relief of pain and fever, or other cold medicines
- (3) Children under 15 years of age
- (4) Pregnant women whose due date is within 12 weeks

#### 2. Do not take any of the following drugs while taking this drug.

Other pain reliever and fever reducers, cold medicines, sedatives or motion-sickness drugs

#### 3. Do not drive or operate machinery after taking this drug.

(Sleepiness, etc. may occur.)

#### 4. Do not drink alcohol before and after taking this drug.

#### 5. Do not use this drug for a prolonged period of time.



®: Trademark

## WHEN to consult a doctor

**1. The following people should consult a doctor, dentist, pharmacist or registered salesperson before taking this drug.**

- (1) Those who are under treatment by a doctor or dentist
- (2) Pregnant women or those who may become pregnant
- (3) Breastfeeding women
- (4) Elderly people
- (5) Those with any previous allergic reactions to drugs or other things
- (6) Those who were diagnosed with any of the following diseases: Heart disease, kidney disease, liver disease, systemic lupus erythematosus or mixed connective-tissue disease
- (7) Those who have suffered from any of the following diseases:  
Gastric and duodenal ulcers, ulcerative colitis or Crohn's disease

**2. The following symptoms may be adverse drug reactions. If any of these symptoms occur, STOP TAKING the drug immediately and consult a doctor, dentist, pharmacist or registered salesperson. Show them this leaflet.**

| Related areas          | Symptoms   |
|------------------------|--|
| Skin                   | Rash, redness, itching or purpura  |
| Digestive system       | Nausea, vomiting, loss of appetite, stomach discomfort, stomachache, oral ulcer, heartburn, heavy stomach feeling, gastrointestinal bleeding, abdominal pain, diarrhea or bloody stool |
| Central nervous system | Dizziness/vertigo  |
| Circulatory system     | Palpitations   |
| Respiratory system     | Breath shortness   |
| Other                  | Blurred vision, ringing in the ears, swelling, nose bleeding, gum bleeding, uncontrolled bleeding, bleeding, back pain, excessive fall in temperature or malaise                       |

In rare cases, the following serious symptoms could occur. In such an event, immediately consult a doctor.

| Symptom's name   | Symptoms   |
|--|--|
| Shock (Anaphylaxis)  | Itching of skin, hives, hoarseness, sneezing, itching of throat, tightness in the chest, palpitations, consciousness clouding, etc. which develop soon after taking the drug |
| Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), Toxic epidermal necrolysis | Persistent or acute worsening of high fever, bloodshot eyes, eye discharge, lip erosion, sore throat, widespread rash, redness, etc.   |

(Continued on the next page)

(Continued from the previous page)

| Symptom's name     | Symptoms  |
|--------------------|---|
| Liver dysfunction  | Fever, itching, rash, jaundice (skin and whites of the eyes turn yellow), brown urine, general malaise, loss of appetite, etc.  |
| Kidney disorders   | Fever, rash, reduction of urine output, generalized swelling, general malaise, joint pain (painful joints), diarrhea, etc.  |
| Aseptic meningitis | Severe headache with neck muscle rigidity, fever, nausea, vomiting, etc. (These symptoms are often reported mainly in patients undergoing treatment for systemic lupus erythematosus or mixed connective-tissue disease). |
| Asthma             | Wheezing or crowing sound when breathing breathlessness, etc.   |
| Aplastic anemia    | Purpura, nose bleeding, gum bleeding, fever, pale skin, pale mucosa, tiredness, palpitations, breathlessness, giddy feeling accompanied by queasy feeling, bloody urine, etc.   |
| Agranulocytosis    | Sudden high fever, chills, sore throat, etc.  |

**3. You may develop the following symptoms after taking this drug. If the symptoms continue or worsen, stop taking the drug and consult a doctor, pharmacist, or registered salesperson. Show them this leaflet.**

Constipation, sleepiness

**4. If the symptoms do not improve after 5 or 6 doses, stop taking this drug and consult a doctor, dentist, pharmacist or registered salesperson. Show them this leaflet.**

## Indications

- Relief of pain associated with headache, stiff-shoulders, toothache, menstruation, sore throat, joints, muscles, neuralgia, lower-back, tooth extraction, bruising, earache, bone fracture, sprains and wounds
- Fever reduction, relief from chills due to fever

## Dosage and Directions

Take the following dose up to 3 times a day, at intervals of at least 4 hours, avoid taking on an empty stomach. This drug should be taken with either cold or lukewarm water.

| Age      | Adult (15 years or over) | Under 15 years |
|----------|--------------------------|----------------|
| One dose | 2 tablets                | Do not take    |

## <Cautions regarding dosage and directions>

(1) Strictly observe the dosage and directions.

(2) <How to remove the tablet>

As shown in the right figure, with the fingertip strongly push the convex portion of the PTP sheet containing a tablet, break the aluminum foil on the back side and take it out.

(Swallowing a tablet without removing the PTP sheet by mistake may cause unexpected problems such as the PTP sheet sticking to the esophageal mucous membrane.)

(How to remove medicine  
from the PTP sheet)



## Ingredients

Two tablets contain:

Ibuprofen . . . . . 150 mg

Magnesium Oxide . . . . . 100mg

Allylisopropylacetylurea . . . . . 60 mg

Anhydrous Caffeine . . . . . 80 mg

## Additives

Anhydrous silicic acid, cellulose, hydroxypropyl cellulose, hypromellose, macrogol, magnesium stearate, talc and titanium oxide

## Storage and Handling Cautions

(1) Store this drug in a cool, dry place away from direct sunlight and moisture.

(2) Keep out of the reach of children.

(3) Do not transfer this drug to a different container (It may cause misuse or deterioration of quality).

(4) Do not use this drug after the expiration date.

製造販売元



**Eisai製薬株式会社**

〒163-1488 東京都新宿区西新宿3-20-2  
<https://www.ssp.co.jp/>

The marketing of this drug has been approved for domestic sales in accordance with the Japanese Pharmaceutical and Medical Devices Act (PMD Act.)